

## SHORT COMMUNICATION

**Screening for Resistance against Northern Corn Leaf Blight (*Exserohilum turcicum* (Pass.) K.J. Leonard & Suggs) in Temperate Maize Lines****Babita Chaudhary<sup>1</sup> and VP Mani<sup>2</sup>**<sup>1</sup>*Division of Genetics, IARI, New Delhi-110012*<sup>2</sup>*Vivekananda Parvatiya Krishi Anusandhan Sansthan, Almora-263 601, Uttarakhand*

(Received 14 December 2009; Revised 20 June 2011; Accepted 25 July 2011)

Ten early duration maize lines, namely, CM 128, V 327, V 335, V 13, V 128, V 17, V 336, CM 129, V 53 and V 27, were screened against northern corn leaf blight (*Exserohilum turcicum* (Pass.) K.J. Leonard & Suggs) under field as well as under artificial epiphytotic conditions. Five inbreds, viz., V 335, V 13, V 336, V 53 and V 27 were identified as resistant while rest five viz., CM 128, V 128, V 327, V 17 and CM 129 were found susceptible. In the pooled disease scores across the three seasons, the highest mean score of 4.0 was recorded for V 128, indicating high level of susceptibility, whereas, lowest mean score of 1.5 recorded for V 335, showed high level of resistance across the locations. The promising inbreds can be utilized as a resistant donor in breeding programme. These inbreds possess resistance to northern corn leaf blight, can be used successfully in developing high yielding early maturing varieties for the hill region.

**Key Words:** *Exserohilum turcicum*, Maize, Resistance, Screening, Turcicum leaf blight

The production of the maize in hilly areas (Kumaoun hills) of India is low as compared to the other area of the nation. The major cause of low yield is Northern corn leaf blight which causes a wide range (28-91%) of yield losses in maize (Sharma and Aujla, 1968; Pant *et al.*, 2000; Singh *et al.*, 2003).

Success of any crop improvement programme depends on the nature and magnitude of genetic variability present in the crop. The effectiveness of selection largely depends on the amount of heritability for traits under study. Therefore, present study was undertaken to reduce the loss due to Turcicum leaf blight (*Helminthosporium turcicum* Pass. or *Exserohilum turcicum* (Pass.) K.J. Leonard & Suggs) through selecting early maturing resistant maize material. An attempt has been made to evaluate the variability in present material of maize.

Turcicum leaf blight is the most common as well as chronic problem of maize in the hills. Passerini, from Italy was first to describe the Turcicum leaf blight in 1876. In India, Butler *et al.* (1920), first observed this disease in maize. It is the most prevalent form in all the major maize growing regions of India during rainy (*kharif*) as well as winter (*rabi*) season (Lal, 1991). The disease can substantially reduce the grain

yield of maize over a wide range from 28% to 91% (Ullstrup, 1951; Chenulu and Hora, 1962; Sharma and Aujla, 1968; Pant *et al.*, 2000). Severe disease conditions prevailing at the time of ear formation may cause total grain loss.

This disease has also been observed to be the most important in the hills, mainly in Jammu and Kashmir, Himachal Pradesh, Sikkim, West Bengal, Meghalaya, Tripura, Assam and Uttarakhand. The symptoms first develop on lower leaves and successively progress to upper leaves of the plant under favourable conditions. In severe disease condition, the entire foliage generally gets killed, giving the appearance of frosted leaves. The ears and kernels are seldom infected, but some lesions do develop in the outer husk. The typical symptoms seen on a susceptible host are long elliptical (4 to 20 cm long and 1 to 5 cm wide), grayish-green to brown (tan) in color. Spores are produced in the lesions, which are olive-green to black, and usually produce concentric rings, giving the entire spot a target like appearance. Spores from the primary lesions re-infect the host producing secondary cycle of the disease (Partridge, 1997). The presence and absence of this type of lesions are governed by host-pathogen interaction (Carlos De Leon, 1987).

\* Author for Correspondence E-mail: babchaudhary@yahoo.co.in

Ten maize inbred lines with different genetic background and origin were evaluated against Turcicum leaf blight under artificial epiphytotic conditions during *kharif* 1998 and 1999 at Hawalbagh, Almora and *rabi* 1998-99 at Amberpet, Hyderabad. Heavily infected leaves with Turcicum leaf blight collected during last season and were ground to powder and stored at room temperature. The powdered inoculum was inoculated as per procedure. A pinch of the leaf powder was dropped into the whorl of each plant, commencing from 20 days after planting. The inoculation was preferably done late in the afternoon till the pre-tasseling stage four times at weekly intervals. Spray of spore suspension with spore concentration of  $2.5 \times 10^4/\text{ml}$  was done on plants, commencing from 20 days after planting till pre-tasseling stage at 5-7 days interval. The disease severity was recorded at flowering stage and dry silk stage in terms of intensity following the rating scale of 1.0 to 5.0 with 0.5 intermediate rating (Payak and Sharma, 1982). Intermediate ratings between two numbers (1.25, 1.5, 1.75, 2.25, 2.75, 3.25, 3.5, 3.75, 4.25, 4.5 and 4.75) was also recorded. Row-to-row spacing of 60 cm and plant-to-plant of 25 cm was maintained at both the locations. Fertilizer doses @ of 160 Kg N, 60 Kg  $\text{P}_2\text{O}_5$  and 40 Kg  $\text{K}_2\text{O}$  were applied to the crop at both the locations.

Mean score for Turcicum leaf blight recorded during *kharif* 1998 and 1999 are presented in Table 1. Mean score of turcicum leaf blight during *kharif* 1998 ranged from 1.8 (V 53) to 4.0 (V 17) while, during *kharif* 1999 it varied from 1.9 (V 13) to 4.5 (V 17). The five inbreds, viz., CM 128, V 327, V 128, CM 129 and V 17 exhibited above 3.0 score and were identified as susceptible while rest of five inbreds, V 335, V 13, V 336, V 53 and V 27 with disease of score of less than 2.5, were found to possess resistance. V 17 recorded the maximum disease score of 4.5 during *kharif* 1999, was highly susceptible, whereas the minimum disease score of 1.8 was recorded for V 53 during *kharif* 1998, which indicated high level of resistance for Turcicum leaf blight.

The disease mean scores for Turcicum leaf blight recorded during *rabi* 1998-1999 are presented in Table 2. The highest mean score (3.9) was recorded for V 17, while the lowest (1.5) for V 335. Out of ten inbreds tested during the season, five, viz., V 335, V13, V 336, V 53 and V 27 indicated resistance

**Table 1. Mean disease reaction of maize inbreds Turcicum leaf blight, *kharif* 1998 and 1999 at Hawalbagh, Almora**

Inbred lines	1998	1999	Mean	Type of reaction
CM 128	3.5	3.8	3.7	S
V 327	3.8	4.1	3.9	S
V 335	2.0	2.0	2.0	R
V 13	2.3	1.9	2.1	R
V 128	3.8	4.1	3.9	S
V 17	4.0	4.5	4.3	S
V 336	2.0	2.3	2.1	R
CM 129	3.0	3.5	3.3	S
V 53	1.8	2.0	1.9	R
V 27	2.3	2.5	2.4	R

S-susceptible (2.6 to 5.0), R-resistant (1.0 to 2.5)

**Table 2. Mean disease reaction of maize inbreds for Turcicum leaf blight, *rabi* 1998-99 at Amberpet, Hyderabad**

Inbred lines	1998-99	Type of reaction
CM 128	3.0	S
V 327	3.3	S
V 335	1.5	R
V 13	1.7	R
V 128	4.0	S
V 17	3.9	S
V 336	2.0	R
CM 129	3.5	S
V 53	2.2	R
V 27	2.0	R

S-susceptible (2.6 to 5.0), R-resistant (1.0 to 2.5)

**Table 3. Pooled mean disease reaction of maize inbreds Turcicum leaf blight (Hawalbagh and Hyderabad)**

Inbred lines	1998 ( <i>Kharif</i> )	1998-99 ( <i>Rabi</i> )	1999 ( <i>Kharif</i> )	Pooled	Type of reaction
CM 128	3.5	3.0	3.8	3.4	S
V 327	3.8	3.3	4.1	3.7	S
V 335	2.0	1.5	2.0	1.8	R
V 13	2.3	1.7	2.0	2.0	R
V 128	3.8	4.0	4.1	4.0	S
V 17	4.0	3.9	4.5	4.1	S
V 336	2.0	2.0	2.3	2.1	R
CM 129	3.0	3.5	3.5	3.3	S
V 53	1.8	2.1	2.0	2.0	R
V 27	2.3	2.0	2.5	2.3	R

S-susceptible (2.6 to 5.0), R-resistant (1.0 to 2.5)

for Turcicum leaf blight, while remaining five (CM 128, V 327, V 128, V 17 and CM 129) showed susceptibility.

The disease reaction for Turcicum leaf blight was pooled across three seasons in two environments and is presented in the Table 3. The highest mean score

of 4.0 was recorded for V 128, indicating highest level of susceptibility and lowest mean score (1.8) was recorded for V 335 indicating highest level of resistance across the environment. The other susceptible inbred lines on the basis of pooled score were, CM 128 (4.0), V 327 (3.7), V 128 (4.0), V 17 (4.1) and CM 129 (3.3), while inbreds having score of 1.8 (V 335), 2.0 (V 13), 2.1 (V 336), 2.0 (V 53) and 2.3 (V 27) were resistant to Turcicum leaf blight.

In a similar study, Singh *et al.* (2003), identified 17, 27, 111 and 10 inbred lines developed at Almora as resistant, moderately resistant, susceptible and highly susceptible to Maydis leaf blight caused by *Bipolaris maydis* (*Cochliobolus heterostrophus*). Their study, which involved about 65 genotypes, also indicated susceptibility of CM 128, V 327, V 128, V 17 and V 27, while V 335, V 13, V 336 and V 53 showed resistance reaction.

Elliott and Jenkins (1946) screened 200 inbred lines, 126 crosses and 184 double crosses against the Turcicum leaf blight and found that NC34 was the most resistant while CI123, K715, KY114, MO21A, T49B, T105, BK115, CI15 and TX116 showed traces of infection and the resistance to Turcicum leaf blight was transmitted to hybrid progeny. In a similar study, Gowda and Kaiser (1990) tested 37 maize lines for resistance to *Exserohilum turcicum*. Of these 10 were resistant and 12 were moderately resistant. Rathee *et al.* (1999) reported eight genotypes having high level of disease resistance.

The inbred lines identified to possess resistance to Turcicum leaf blight in the present study, can be used successfully in developing high yielding early maturing varieties for the hill region having high level of resistance to Turcicum leaf blight.

## References

- Butler EJF, JF Shaw and AK Mitra (1920) Reports of imperial mycologist. *Reports of Agricultural Research Institute and College*, Pusa, New Delhi.
- Carlos De leon (1987) *Exserohilum turcicum* blight of corn, the blighter and the blighted. Paper presented at the XXX National Maize Workshop, March 17-20, 1987, Hyderabad, Andhra Pradesh, India.
- Chenulu VV and Hora TS (1962) Studies on losses due to *Helminthosporium* blight of maize. *Indian Phytopath.* **15**: 235.
- Elliott C and MT Jenkins (1946) *Helminthosporium turcicum* leaf blight of corn. *Phytopathology* **36**: 660-666.
- Gowda KTP and SAKM Kaiser (1990) Evaluation of some elite inbred lines of maize for resistance to turcicum leaf blight in Karnataka, India and their prospects on hybrid development. *J. Mycopath. Res.* **28**: 75-80.
- Lal S (1991) Genetics of *Helminthosporium* leaf blight resistance in maize. In: KR Sarkar, NN Singh and JKS Sachan (eds.) *Maize Genetics Perspectives, Indian Society of Genetics and Plant Breeding*, New Delhi, pp 231-237.
- Pant SK, Kumar Pramod and VS Chauhan (2000) Effect of turcicum leaf blight on photosynthesis in maize. *Indian Phytopath.* **54**: 251-252.
- Payak MM and RC Sharma (1982) Disease rating scales in maize in India. Techniques of scoring for resistance to important diseases of maize. *All India Co-ordinated Maize Improvement Programme*, ICAR, New Delhi.
- Partridge JE (1997) *Northern Corn Leaf Blight*. University of Nebraska – Lincoln, USA.
- Rathee VK, BK Sharma and SK Guleria (1999) Screening of maize germplasm against turcicum leaf blight under artificial epiphytotic conditions. *Crop Res. Hissar* **17**: 268-269.
- Singh R, VP Mani, RS Khendelwal, P Bhandari, GS Bisht and KS Koranga (2004) Screening of maize genotypes against foliar disease Southern corn leaf blight. *SABRAO J. Breed. Genet.* **36**: 45-47.
- Sharma D. and SS Aujla (1968) Effect of leaf blight on susceptible and resistant stock of maize in Kullu Valley. *J. Res. Punjab Agric. Univ., Ludhiana* **5**: 501-504.
- Ullstrup AJ (1951) The effect of some leaf diseases on grain yield in corn. *Phytopathology* **41**: 36.