

Evaluation and Utilization of Introduced *Musa* Germplasm in India

S Uma, S Sathiamoorthy, MS Saraswathi, P Durai, Hari Om Sharma¹ and Anuradha Agrawal²

¹ National Research Centre for Banana, Tiruchirapalli, Tamil Nadu

² Tissue Culture and Cryopreservation Unit, National Bureau of Plant Genetic Resources, Pusa Campus, New Delhi-110 012

Key Words: Banana, Evaluation, Germplasm, Utilization

Banana and plantains (*Musa* spp.) have got an important niche as a major staple food in India as well as other tropical countries. Their cultivation has become a source of livelihood for the marginal and small farmers across the world. In India, banana and plantains lie interwoven with culture and tradition. India boasts of its polyclonal germplasm with more than 20 varieties contributing to commercial cultivation and over 30 serving as choice cultivars (Uma *et al.*, 2002). But almost all varieties are threatened by a wide range of pests and diseases, and constitute the most significant constraint in *Musa* production. Important diseases of *Musa* include leaf spots diseases (Sigatoka complex) caused by species of *Mycosphaerella*, Fusarium wilt (Panama disease) caused by *Fusarium oxysporum* f. sp. *cubense*, bacterial wilts caused by *Erwinia carotovora* etc. Viral diseases like Banana bunchy top, Banana bract mosaic, Banana streak and Cucumber mosaic are economically important. Pests include nematodes (*Radopholus similis*, *Pratylenchus coffeae*, *Meloidogyne incognita* are important) and the borers (*Cosmopolites sordidus* and *Odoiporus longicollis*). Breeding for resistant cultivars and integrated pest and disease management are the only options for sustainable and economically viable banana production.

Banana is one of the toughest crops for improvement through breeding, due to inherent sterility. However, genetic improvement of diploids (by crossing and selection) started in the early 1930s on the principle that with a generally fixed female parental contribution, success of hybrids must depend totally on the positive genetic qualities transmitted by the pollen. Subsequently tetraploid hybrids were developed; Currently efforts are underway to breed new hybrids through classical breeding as well as biotechnological methods and to popularize them among growers. Various national and international breeding programmes like Fundación Hondureña de Investigación Agrícola (FHIA), Honduras, International Institute for Tropical Agriculture (IITA), Nigeria, Centre de Recherches Regionales sur Bananiers et Plantains

(CRBP), Cameroon, etc. have resulted in the release of more than 25 superior hybrids.

The International Network for Improvement of Banana and Plantains (INIBAP), Montpellier, France, is instrumental in developing a network for collection, conservation and distribution of germplasm and improved global hybrids. Additionally, an International *Musa* Testing Programme (IMTP) is a collaborative effort in which elite *Musa* varieties produced by breeding programmes, as well as promising germplasm accessions from the INIBAP collection, are evaluated in sites worldwide (Orjeda, 2000). IMTP has been running since 1990 as a partnership between National Agriculture Research Systems, INIBAP and breeding programmes. India is a partner in the IMTP in which many hybrids were evaluated. With the National Research Centre for Banana (NRCB), Trichy, Tamil Nadu, as the nodal center, India has four test centers to undertake trials of evaluation of global hybrids against *Fusarium* wilt, Sigatoka, nematodes, and performance in farmers' fields. This paper briefly describes the status of introduced hybrids and germplasm of *Musa* and their extent of utilization in India.

Exotic banana hybrids and cultivars were obtained from the world collection of banana held at INIBAP Transit Centre (ITC), Leuven, Belgium in the form of proliferating *in vitro* cultures. The cultures were quarantined at National Bureau of Plant Genetic Resources (NBPGR), New Delhi and conserved in its *in Vitro* Genebank. They were subcultured in Murashige and Skoog's (1962) medium with 3% sucrose, 1-3 mg/l N⁶-benzyladenine, 0.1-0.5mg/l IAA for bulking up. In a phased manner during 1994-2004, several hybrids (Table 1) and cultivars were supplied to the NRCB for field evaluation and trial in the farmers' fields.

During the second phase of IMTP, 21 accessions were screened against *Fusarium* wilt (*Fusarium oxysporum* race 4) and the results showed that the introductions namely Bluggoe, Burro Cemsa (ITC 1259) and Saba (ITC 1138), despite their greater rating for vascular

Table 1. Details of some of the promising global hybrids evaluated under the IMTP by NRCB, Trichy

INIBAP No.	NBPGR No.	Name of Hybrid	Parentage	Important Traits
<i>Source Institute: International Institute for Tropical Agriculture (IITA), Nigeria</i>				
ITC 1296	EC468354	BITA-2 (TMB x 1378)	Pisang Awak cv. Fougamou X <i>M. balbisiana</i> 1-63	Sturdy, high yielding with resistance to Black Sigatoka
ITC 1297	EC426459	BITA-3 (TMB x 5295)	AAB Laknau, cv. Laknau x Tjau lagada	High yielding with resistance to Black Sigatoka, suitable for culinary purpose
<i>Source Institute: Centre de Recherches Regionales sur Bananiers et Plantains (CRBP), Cameroon</i>				
ITC 1344	EC497313	CRBP-39	AAB – Plantain cv. French clair x M53 (4X)	Highly resistant to Black Sigatoka, possess good cooking qualities
<i>Source Institute: Fundación Hondureña de Investigación Agrícola (FHIA), Honduras</i>				
ITC 0504	EC501358	FHIA-01	AAB – Pome cv. Prata Ana X Prata Ana (Dwarf Prata) SH 3142	Resistant to Black Sigatoka, burrowing nematodes and race 1 and 4 of <i>Fusarium</i> wilt; tolerant to drought and cold, fruits have good apple flavour and long green life; suitable for cooking and beer making
ITC 0505	EC501359	FHIA-02	AAA – Cavendish cv. Williams x SH 3393	Fruits are sweet in taste similar to Cavendish bananas but possess a shorter green life. Susceptible to <i>Fusarium</i> wilt
ITC 0506	EC501360	FHIA-03	(Gaddatu x BB) x SH 2471- SH 3386 SH 3386 x SH 3320	Possess good resistance to Black Sigatoka, tolerant to marginal soil fertility and rainfall, performs well at both sea level and high altitudes; resistant to strong winds, drought and tolerant to cold; hybrid is highly suitable for homestead gardens
ITC 1264	EC407587	FHIA-17	Cultivar AAA – Gros Michel Highgate x SH 3362	Possess good resistance to Black Sigatoka and race 1 of <i>Fusarium</i> wilt
ITC 1319	EC467908	FHIA-18	AAB – Pome cv. Prata Ana x SH 3142	Resistant to Black Sigatoka; fruits are sweet acid in taste with apple flavour
ITC 1332	EC469509	FHIA-21	AAB – Plantain cv. AVP – 67 x SH 3142	French plantain is resistant to Black Sigatoka and has excellent plantain flavour; twice as productive as False Horn; recommended for home gardens
ITC 1265	EC468351	FHIA-23	cv. AAA – Gros Michel Highgate x SH 3362	Tolerant to Black Sigatoka and resistant to race 1 of <i>Fusarium</i> wilt
ITC 1418	EC467911	FHIA-25	SH 3648 (4X) x SH 3142 (2X)	Dwarf and sturdy with high level of resistance to Black Sigatoka; fruits are easy to peel, fast to cook and possess excellent texture and flavour both boiled and fried and long green life after harvest

discolouration performed well and yielded good bunches indicating their resistance to *Fusarium* wilt. Similar screening was done for Sigatoka leaf spot diseases (*Mycosphaerella* spp.) during the third phase of IMTP. Among the nine accessions tested, the exotic hybrids namely TMB 5295-1, FHIA-18, CRBP-39 and FHIA-21 were found to be resistant to Sigatoka leaf spot diseases. When the introductions were screened against the three major nematodes namely *Pratylenchus coffeae*, *Meloidogyne incognita* and *Helicotylenchus multicinctus*, none were found to be resistant or immune to all the three nematodes. However GCTCV 215 was found to be resistant to *P. coffeae* alone while Pisang Berlin (ITC 0611), Pisang Ceylon (ITC 1441) and FHIA-3 were found to be resistant to *H. multicinctus* alone.

Superior hybrids and clones were evaluated at NRCB, Trichy and the promising ones were identified and distributed for multilocation trials under the All

India Coordinated Research Project (AICRP)-Tropical Fruits for commercial cultivation (Anon., 2004). The varieties evaluated are detailed in Table 2.

Out of these varieties, FHIA-01, FHIA-03, FHIA-23 and Saba, which were found to be highly promising, have been taken to farmer's fields for large scale testing under various agro-climatic conditions across the country. The programme on evaluation of global hybrids in the farmers' fields is being partially supported by BAPNET/INIBAP, France. Under this three major hybrids namely FHIA-01, FHIA-03 and FHIA-23 have been multiplied and supplied by NRCB, Trichy to various farmers. FHIA-01 was widely tested in Maharashtra by a leading tissue culture industry and the fruits were evaluated for post-harvest processing and preparation of puree. FHIA-01 was found to be more suitable for processing with high sugar:acid ratio and low polyphenol oxidation of the pulp, which results in pulp browning. The consumers

Table 2. Promising exotic hybrids of *Musa* identified by NRCB for multi-location trials for commercial cultivation under the AICRP-Tropical Fruits

AICRP Centre	Hybrid No.
National Research Centre for Banana, Trichy, Tamil Nadu	FHIA-01, FHIA-03, FHIA-18, FHIA-23, FHIA-21, FHIA-25, CRPB-39, SH-3640
University of Agricultural Sciences, Arabhavi, Dharwad, Karnataka	FHIA-01, FHIA-03, FHIA-17
Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu	FHIA-01, FHIA-03, FHIA-17, FHIA-23
Kerala Agricultural University, Kannara, Kerala	FHIA-17, FHIA-18, FHIA-21, FHIA-25, CRPB-39
Acharya NG Ranga Agricultural University, Kovvur, Andhra Pradesh	FHIA-01, FHIA-03, FHIA-17
Bidhan Chandra Krishi Vishwavidyalaya, Mohanpur, West Bengal	FHIA-01, FHIA-03, FHIA-17
Rajendra Agricultural University, Pusa, Bihar	FHIA-01, FHIA-03, FHIA-17, FHIA-23

have popularly accepted FHIA-03 as cooking banana, as also the cultivar Saba. FHIA-23 is still being evaluated as a dessert cultivar, as substitute for local Cavendish clones.

In the evaluation trial conducted at NRCB, Trichy to assess the superiority of FHIA-01 over the indigenous hybrids and cultivars belonging to the Pome subgroup namely, H-1 and Pachanadan, FHIA-01 was found to be promising for many traits. Tolerance to wind damage was attributed to plant robustness, high yield, extended green and yellow shelf life, resistance to *Fusarium* and Sigatoka and tolerance to weevils.

References

- Anonymous (2004). *AICRP-Tropical Fruits – Research Report*. Tech. Doc. No.80., 326p.
- Murashige T and F Skoog (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* **15**: 473-497.
- Orjeda G (2000) IMTP – A worldwide effort of *Musa* scientific community. In: HP Singh and KL Chadha (Eds.) *Banana – Improvement, Production and Utilisation*. AIPUB, National Research Centre for Banana, Tiruchirapalli, Tamil Nadu, p. 6-10.
- Uma S and S Sathiamoorthy (2002) *Names and Synonyms of Bananas and Plantains in India*. National Research Centre for Banana, Tiruchirapalli, Tamil Nadu, 60p.

Status of Introduced *Musa* Germplasm at Banana Research Station, Kannara

Rema Menon¹, Anuradha Agrawal² and BMC Reddy¹

Banana Research Station, Kannara, Thrissur-680 652, Kerala

¹ *Indian Institute of Horticultural Research, Hessaraghatta, Bangalore, Karnataka*

² *National Bureau of Plant Genetic Resources, Pusa Campus, New Delhi-110 012*

Key Words: *Musa*, Germplasm, Conservation, Genepool, Cultivars, Wild Species

Musa germplasm conservation and evaluation activities are in progress under the All India Coordinated Research Project on Tropical Fruits at the Banana Research Station (BRS), Kannara. The work aims to collect and conserve wild and cultivated germplasm, representative of the diversity existing in the *Musa* genepool for evaluation and subsequent use in breeding programme and to make available selected superior material to the farmers of Kerala. The center presently holds a germplasm of 256 accessions in its field genebank comprising indigenous and exotic varieties.

Since 1994, efforts have been made to enrich the germplasm collection through introductions and till 2004, 48 accessions have been received. Exotic germplasm

was procured by the National Bureau of Plant Genetic Resources (NBPGR), New Delhi, from the global *Musa* germplasm collection held at International Network for Improvement of Banana and Plantain (INIBAP) at its Transit Centre (ITC) in Leuven, Belgium.

Materials and Methods

Proliferating *in vitro* cultures of *Musa*, received from INIBAP, were multiplied at NBPGR, New Delhi and periodically supplied to BRS, Kannara during 1994-2004. The 48 accessions comprising 21 germplasm/reference cultivars, 6 wild species and 21 improved hybrids/variants were received in the form of small suckers or proliferating *in vitro* cultures which were regenerated to plantlets at the Center for Plant